



Nez Perce County

215 10th Street Lewiston, ID 83501 (208) 799-3100 Fax (208) 799-0349

Latah County

333 E Palouse River Drive Moscow, ID 83843 (208) 882-7506 Fax (208) 882-3494

Clearwater County

105 115th Street Orofino, ID 83544 (208) 476-7850 Fax (208) 476-7494

Idaho County

903 West Main Grangeville, ID 83530 (208) 983-2842 Fax (208) 983-2845

Lewis County

132 N Hill Street P O Box 277 Kamiah, ID 83536 (208) 935-2124 Fax (208) 935-0223

Physician's ALERT...

January 23, 2020

Interim Guidance for Healthcare Providers on 2019 Novel Coronavirus (2019-nCoV)

This is an emerging, rapidly evolving situation and updated information will be provided as it becomes available.

Summary

Health authorities in China continue to investigate an outbreak of novel coronavirus (2019-nCoV) infections associated with a large live animal market in Wuhan City, Hubei Province, China. Although person-to-person transmission has been indicated, it is not currently known how easily it is spread person-to-person. Since December 2019, confirmed cases of 2019-nCoV have occurred in an increasing number of countries. Several deaths have been reported among about 300 reported cases. The first confirmed case of 2019-nCoV in the United States was announced, in a Washington State resident recently returned from travel to Wuhan City. On 1/17/2020, screening of travelers from Wuhan City began at three major US airports

Recommendations for Healthcare Providers

- > Obtain detailed travel history on patients with fever and acute respiratory illness.
- Report "patients under investigation" (PUI) for 2019-nCoV infection (see below) to your local public health district or the Bureau of Communicable Disease Prevention (208-334-5939) immediately.

Criteria for a PUI in association with the outbreak of 2019-nCoV in Wuhan City, China are:

- 1) Fever AND symptoms of lower respiratory illness (e.g., cough, shortness of breath)
 - and in the last 14 days before symptom onset,
 - History of travel from Wuhan City, China -or-
 - Close contact² with a person who is under investigation for 2019-nCOV while that person was ill.
- 2) Fever OR symptoms of lower respiratory illness (e.g., cough, shortness of breath)
 - and in the last 14 days before symptom onset,
 - Close contact² with an ill laboratory-confirmed 2019-nCoV patient.

Criteria may be updated at https://www.cdc.gov/coronavirus/novel-coronavirus-2019/clinical-criteria.html. Contact your local public health district or the Bureau of Communicable Disease Prevention if you have questions about clinical presentation, travel history, or exposure criteria.

Notify infection control personnel in your healthcare facility <u>immediately</u> if you identify a PUI for 2019-nCoV.

Collect specimens for 2019-nCoV testing.

Collect lower respiratory, upper respiratory, and serum specimens as soon as possible once a PUI is identified regardless when symptoms began. Additional specimen types (e.g., stool, urine) may be collected and stored. Submit specimens to the Idaho Bureau of Laboratories (IBL) for 2019-nCoV testing at the Centers for Disease Control and Prevention. Do not wait for results on testing for other respiratory pathogens. Contact IBL at 208-334-2235 before submitting specimens. Contact IBL or see https://www.cdc.gov/coronavirus/2019-nCoV for additional guidance.

Clinical laboratories should not attempt virus isolation in cell culture or initial characterization of viral agents recovered in cultures of specimens from a PUI for 2019-nCoV.

> Take precautions.

PUI should wear a surgical mask as soon as they are identified. Evaluate PUI in a private room with the door closed, ideally an airborne infection isolation room. Healthcare personnel entering the room should use standard precautions, contact precautions, airborne precautions, and use eye protection (e.g., goggles or a face shield). See https://www.cdc.gov/coronavirus/2019-nCoV/infection-control.html for infection prevention and control updates.

See https://.cdc.gov/coronavirus/2019-ncov/index.html for more information as the situation evolves.

Footnotes

¹Fever may not be present in some patients, such as those who are very young, elderly, immunosuppressed, or taking certain fever-lowering medications. Clinical judgment should be used to guide testing of patients in such situations.

- ² Data to inform the definition of close contact are limited. Close contact with a PUI for 2019-nCOV is defined as—
- a) being within approximately 6 feet (2 meters), or within the room or care area, of a novel coronavirus case for a prolonged period of time while not wearing recommended personal protective equipment or PPE (e.g., gowns, gloves, NIOSH-certified disposable N95 respirator, eye protection); close contact can include caring for, living with, visiting, or sharing a healthcare waiting area or room with a novel coronavirus case.

- or -

b) having direct contact with infectious secretions of a novel coronavirus case (e.g., being coughed on) while not wearing recommended personal protective equipment.

Considerations when assessing close contact include the duration of exposure (e.g., longer exposure time likely increases exposure risk) and the clinical symptoms of the person with novel coronavirus (e.g., coughing likely increases exposure risk as does exposure to a severely ill patient). Special consideration should be given to those exposed in healthcare settings.

For additional information or to report a case, please contact

Anna Olson RN, BSN, Epidemiologist
Public Health - Idaho North Central District, (208) 799-0387
After Hours Contact: State Communications (State Comm) 800-632-8000